arXiv:cond-mat/0511014v1 [cond-mat.other] 1 Nov 2005

Entanglement and alpha entropies for a massive scalar field in two dimensions

H. Casini* and M. Huerta†
Centro Atómico Bariloche, 8400-S.C. de Bariloche, Río Negro, Argentina

Abstract

We find the analytic expression of $\mathrm{tr}\rho^n(L)$ for a massive boson field in 1+1 dimensions, where $\rho(L)$ is the reduced density matrix corresponding to an interval of length L. This is given exactly (except for a non universal factor) in terms of a finite sum of solutions of non linear differential equations of the Painlevé V type. Our method is a generalization of one introduced by Myers and is based on the explicit calculation of quantities related to the Green function on a plane, where boundary conditions are imposed on a finite cut. It is shown that the associated partition function is related to correlators of exponential operators in the Sine-Gordon model in agreement with a result by Delfino et al. We also compute the short and long distance leading terms of the entanglement entropy. We find that the bosonic entropic c-function interpolates between the Dirac and Majorana fermion ones given in a previous paper. Finally, we study some universal terms for the entanglement entropy in arbitrary dimensions which, in the case of free fields, can be expressed in terms of the two dimensional entropy functions.

1 Introduction

The trace of the vacuum state projector over the degrees of freedom lying outside a given set A gives place to a local density matrix ρ_A, which is generally mixed. The associated entropy $S(A)$ is called geometric or entanglement entropy. In recent years there has been much interest on the properties of the local reduced density matrices. This is partially motivated by ideas related to black hole physics (i.e. see [1, 2, 3, 4]), developments in quantum information theory, and the density matrix renormalization group methods in two dimensions (within the area of condensed matter physics see for example [5, 6]). Besides, the entanglement entropy as well as other measures of information for the reduced density matrices of the vacuum state are in its own right interesting and relevant quantities in the context of quantum field theory. For example, they show a different structure of divergences and a nice interplay between geometry and ultraviolet behavior. Among the results in this sense we can mention the existence of an entropic c-theorem [7] and the recent discovery of a topological entanglement entropy [8].

In a previous paper we studied universal quantities derived from the local density matrix ρ_A, for a Dirac field in two dimensions [9]. In the present work we find the corresponding results for a massive scalar field. Specifically, we shall consider the α entropies

$$S_\alpha(A) = \frac{1}{1-\alpha}\log \mathrm{tr}(\rho_A^\alpha)\,, \tag{1}$$

and the entanglement entropy

$$S(A) = -\mathrm{tr}(\rho_A \log \rho_A) = \lim_{\alpha\to 1} S_\alpha(A)\,, \tag{2}$$

where A is a single interval of length L. As these quantities are not well defined in the continuum limit, we work instead with its associated dimensionless and universal functions $c_\alpha(L)$ and $c(L)$

$$c_\alpha(L) \;\equiv\; L\frac{dS_\alpha(L)}{dL}\;\;,\quad c(L) \;\equiv\; c_1(L)\;. \tag{3}$$

*e-mail: casini@cab.cnea.gov.ar
†e-mail: huerta@cabtep2.cnea.gov.ar

The entropic c-function $c(L)$ is always positive and decreasing, and plays the role of Zamolodchikov's c-function in the entanglement entropy c-theorem [7].

The traces $\mathrm{tr}\rho_A^\alpha$ involved in (1), with $\alpha = n \in \mathbb{Z}$, can be represented by a functional integral on an n-sheeted surface with conical singularities located at the boundary points of the set A [5, 9]. Here, calling u and v to the end points of the segment, the replicated space is obtained considering n copies of the plane cut along the interval (u, v), sewing together the upper side of the cut $(u, v)^k_{\rm out}$ with the lower one $(u, v)^{k+1}_{\rm in}$, for the different copies $k = 1, ..., n$, and where the copy $n + 1$ coincides with the first one. The trace of ρ^n is then given by the functional integral $Z[n]$ for the field in this manifold,

$$\mathrm{tr}\rho^n = \frac{Z[n]}{Z[1]^n}\,. \tag{4}$$

We consider a complex scalar field and following [9], we map this problem into the one of a single cut plane with n decoupled fields $\Phi_k(x)$, $k = 0, ..., n - 1$. These are multivalued, since when encircling the points u or v in anti-clockwise sense they get multiplied by $e^{i\frac{k}{n}2\pi}$ or $e^{-i\frac{k}{n}2\pi}$ respectively, with $k = 0, ..., n - 1$. Then

$$\log Z[n] = \sum_{k=0}^{n-1} \log Z_{k/n}\,. \tag{5}$$

where Z_a, with $a \in [0, 1]$, is the partition function of a complex scalar field in a plane with boundary condition

$$\Phi_{\rm in}(x) = e^{i2\pi a}\Phi_{\rm out}(x) \tag{6}$$

along the cut (u, v).

The free energy and the Green function are related by the identity

$$\partial_{m^2} \log Z_a = -\int dr^2 G(\vec{r}, \vec{r})\,. \tag{7}$$

Our strategy is to find the right hand side of this equation. The information on the Green function on a cut plane which is relevant to the present problem is obtained following a generalized version of a method introduced in [10] to deal with Neumann and Dirichlet boundary conditions on the finite cut. It essentially consists in exploiting the rotation and translation symmetry of the Helmholtz equation, even in the presence of the cut boundary conditions, by analyzing the behaviour at the singular points. In Section 2 we adapt this method to our particular boundary conditions and find an exact expression for $\partial_L\partial_{m^2} \log Z_\alpha$ in terms of the solution of a second order non linear differential equation of the Painlevé V type. This equation allow us to identify the partition function with the inverse of a correlator of exponential operators in the Sine-Gordon (SG) model (a related mapping for correlators of boson disorder operators was given in [11]).

This is reminiscent to the fermionic case, where there is also an expression for c_α in terms of SG correlators. This is shown in a previous paper by a very different method involving dualization [9]. The relation between the fermionic and bosonic cases is analyzed in detail in Section 3, where we first present the results for the bosonic α entropies. In this Section we also study the short and long distance expansions for the entanglement entropy, and we show that the entropic c-function for a real scalar interpolates between the ones corresponding to a Dirac and Majorana fields. In the Section 4 we present some results concerning universal terms for the geometric entropy in arbitrary dimensions. These can be exactly computed for free fields in terms of the two dimensional entropic c-functions. The Section 5 contains the conclusions.

2 The Green function

In this section we study the Green function $G(\vec{r}, \vec{r'})$ using a method introduced by Myers in [10]. The following three requirements uniquely define $G(\vec{r}, \vec{r'})$:

a.- It satisfies the Helmholtz equation

$$\left(-\Delta_{\vec{r}} + m^2\right) G(\vec{r}, \vec{r'}) = \delta(\vec{r} - \vec{r'})\,. \tag{8}$$

b.- The boundary conditions are (they also hold for the Green function derivatives)

$$\lim_{\epsilon\to 0^+} G((x,\epsilon),\vec{r}^{\,'}) = e^{i2\pi a}\lim_{\epsilon\to 0^+} G((x,-\epsilon),\vec{r}^{\,'}) \qquad \text{for } x\in[L_2,L_1]\,, \tag{9}$$

$$\lim_{|\vec{r}|\to\infty} G(\vec{r},\vec{r}^{\,'}) = 0\,. \tag{10}$$

Here the interval $[L_2, L_1]$ on the x axis is the cut location. From now on we choose $a \in [0, 1]$.

c.- $G(\vec{r},\vec{r'})$ is bounded everywhere (including the cut) except at $\vec{r}=\vec{r}^{\,'}$.

We will write the Green function as $G(z, z')$ as a shortcut of $G(z, \bar{z}, z', \bar{z}', L_1, L_2)$, where z and $\bar{z}$ are the complex coordinates $x + iy$ and $x - iy$. It is Hermitian

$$G(z,z') = G(z',z)^*\,, \tag{11}$$

and the symmetry of the problem also gives

$$G(z,z')^* = G(\bar{z},\bar{z}'). \tag{12}$$

The reflexion operation

$$R\ (x,y) = (L_1 + L_2 - x, y) \tag{13}$$

leaves the Helmholtz equation, the cut, and boundary conditions invariant. Thus we have

$$G(z,z') = G(Rz, Rz')\,. \tag{14}$$

Due to the boundary conditions, near the end points of the interval $[L_2, L_1]$ the Green function must have branch cut singularities. The requirement that the function must remain bounded on the cut and the equation (8) imply that the most singular terms of $G(z, z')$ for z near L_1 (and fixed z') have to be of the form

$$G(z,z') \sim (z-L_1)^a S_1(z') + (\bar{z}-L_1)^{1-a} S_2(z')\,. \tag{15}$$

We have written explicitly only the terms with powers of $z - L_1$ with exponent smaller than one. These are the ones of interest in the sequel. Note that the contributions at this order must be analytic or anti-analytic in z in order to cancel the Laplacian term in (8).

The most singular contributions to $\partial_{L_1} G(z, z')$ for $z \to L_1$ follow from the derivative of (15)

$$\partial_{L_1} G(z,z') \sim -a(z-L_1)^{a-1} S_1(z') + (a-1)(\bar{z}-L_1)^{-a} S_2(z')\,. \tag{16}$$

The function $\partial_{L_1} G(z, z')$ satisfies the equation (8) and the boundary conditions, and it is not singular at $z \to z'$. It has only one singularity located at L_1 whose expression is given by (16). Therefore the combination

$$\begin{aligned}&\partial_{L_1} G(z,z')(S_1(z'')S_2(z''') - S_1(z''')S_2(z'')) + \partial_{L_1} G(z,z'')(S_1(z''')S_2(z') - S_1(z')S_2(z'''))\\ &+\partial_{L_1} G(z,z''')(S_1(z')S_2(z'') - S_1(z'')S_2(z'))\end{aligned} \tag{17}$$

is nonsingular everywhere, and the uniqueness of the solution implies that it must be identically zero. Thus,

$$\partial_{L_1} G(z,z') = F_1(z)S_1(z') + F_2(z)S_2(z') = F^T(z)S(z') \tag{18}$$

for certain functions $F_1(z)$ and $F_2(z)$. Note that in the last equation we have written the sum over two terms in vector notation. From the Hermiticity relation we have

$$S(z) = A^{-1}\, F(z)^*\,, \tag{19}$$

where A is a Hermitian matrix. Finally we get the following fundamental relation

$$\partial_{L_1} G(z,z') = F^T(z)S(z') = S(z)^\dagger A S(z')\,. \tag{20}$$

The function $S(z)$ satisfies the homogeneous Helmholtz equation with the same boundary conditions as $G(z', z)$ with z' fixed, but it is unbounded around L_1. In fact, it follows from (16) and (20) that it has singular terms proportional to $(z - L_1)^{-a}$ and $(\bar{z} - L_1)^{a-1}$. However, these disappear if we derive with respect to L_2. On the other hand $S(z)$ behaves around L_2 as an ordinary wave, that is, it vanishes proportionally to $(L_2 - z)^a$ and $(L_2 - \bar{z})^{1-a}$. Then $\partial_{L_2} S(z)$ has singular terms around L_2 which are proportional to $(L_2 - z)^{a-1}$ and $(L_2 - \bar{z})^{-a}$. These have the same form as the singular terms of $S(Rz)$ for z near L_2 once one takes into account that

$$z - L_1 = L_2 - R\bar{z}\,. \tag{21}$$

In consequence, there exist a matrix γ such that $\partial_{L_2} S(z) - \gamma\, S(Rz)$ is bounded, and thus identically zero,

$$\partial_{L_2} S(z) = \gamma\, S(Rz)\,, \tag{22}$$

where γ is a function of $L = L_1 - L_2$. Translational invariance then implies

$$\partial_{L_1} S(Rz, L_1, L_2) = (\partial_1 + \partial_2)(S)(Rz, L_1, L_2) = -\partial_3(S)(Rz, L_1, L_2) = -\gamma\, S(z)\,. \tag{23}$$

Using (20), (22) and (23) to calculate $\partial_{L_1}\partial_{L_2} G(z, z') - \partial_{L_2}\partial_{L_1} G(z, z') = 0$ we get that the matrix A must be a constant, $\partial A/\partial_{L_1} = \partial A/\partial_{L_2} = 0$, and it also holds

$$\gamma^\dagger = A\gamma A^{-1}\,. \tag{24}$$

The constant matrix A can then be obtained using the solution for the half infinity cut, which can be computed by standard methods. In polar coordinates the Green function with $L_2 = -\infty$, $L_1 = 0$ is

$$G(r, \theta, r', \theta') = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{i(n+a)\theta} e^{-i(n+a)\theta'} I_{|n+a|}(mr_<) K_{|n+a|}(mr_>)\,, \tag{25}$$

where I and K are the standard modified Bessel functions and $r_>$ and $r_<$ are the maximum and minimum between r and r'. From here we get

$$A = -4\pi(1-a)a\,\sigma_1\,, \tag{26}$$

where σ_1 is the Pauli matrix. The equation (16) leads to the behavior

$$S(z) \sim \frac{1}{4\pi} \begin{pmatrix} \frac{1}{a}(z - L_1)^{-a} \\ \frac{1}{(1-a)}(\bar{z} - L_1)^{a-1} \end{pmatrix} \tag{27}$$

for z in the vicinity of L_1.

2.1 Equations for $S(z)$ from translation and rotation symmetries

In order to use eq. (20) to compute the partition function we need more information on $S(z)$. With this aim we exploit the symmetries of the problem as follows. Translations in the y direction commute with the Helmholtz equation and thus the function $(\partial_y + \partial_{y'})G(z, z')$ is non singular at $z = z'$ and satisfies the equation (8) plus the boundary conditions. However, due to the $\partial_y G(z, z')$ term, it is singular at L_1 and L_2 as a function of z. Near L_1 we have from (15), (19) and (27)

$$\partial_y G(z, z') \sim ia(z - L_1)^{a-1} S_1(z') - i(1-a)(\bar{z} - L_1)^{-a} S_2(z') \sim -iF_1(z)S_1(z') + iF_2(z)S_2(z')\,, \tag{28}$$

and near L_2 we have by the symmetry (14)

$$\partial_y G(z, z') \sim -iF_1(Rz)S_1(Rz') + iF_2(Rz)S_2(Rz'). \tag{29}$$

Then, by subtracting these terms from $(\partial_y + \partial_{y'})G(z, z')$ we obtain a singularity free function which respects the boundary conditions and satisfies the Helmholtz equation. In consequence it is the null function,

$$\partial_y G(z, z') + \partial_{y'} G(z, z') = -i(F^T(z)\sigma_3 S(z') + F^T(Rz)\sigma_3 S(Rz'))\,. \tag{30}$$

Taking the derivative with respect to L_1 we arrive at

$$\begin{aligned}\partial_y F^T(z)S(z') \quad + \quad & F^T(z)\partial_{y'}S(z') = i\left(F^T(Rz)\left\{\gamma,\sigma_3\right\}S(z') + \right. \\ & \left. F^T(z)\left\{\gamma,\sigma_3\right\}S(Rz') + \partial_x F^T(z)\sigma_3 S(z') + F^T(z)\sigma_3\partial_{x'}S(z')\right)\,,\end{aligned} \tag{31}$$

with $\{\gamma,\sigma_3\} = \gamma\sigma_3 + \sigma_3\gamma$. This equation has to be satisfied in z and z' independently. In particular for $z \to L_1$ the coefficients of the singular terms on both sides of (31) must be equal. From (19) and (27) the most singular terms in $\partial_y F^T(z) - i\partial_x F^T(z)\sigma_3$ (which would be proportional to $(z-L_1)^{a-2}$ and $(\bar{z}-L_1)^{-a-1}$) exactly cancel. But this combination of derivatives should also have a contribution to the next orders, $(z-L_1)^{a-1}$ and $(\bar{z}-L_1)^{-a}$, for z near L_1. Then we can write near this point

$$\partial_y F^T(z) - i\partial_x F^T(z)\sigma_3 \sim F^T(z)\xi\,, \tag{32}$$

where ξ is a matrix which depends on L. Thus, isolating the singular terms of (31) for z near L_1, we have

$$\partial_y S(z) = i\left\{\gamma,\sigma_3\right\}S(Rz) + i\sigma_3\partial_x S(z) - \xi S(z)\,. \tag{33}$$

Replacing this back into (31) and using $\{A,\sigma_3\} = 0$, $\{\gamma,\sigma_3\}^\dagger A + A\left\{\gamma,\sigma_3\right\} = 0$, we obtain

$$\xi^\dagger A + A\xi = 0\,. \tag{34}$$

We can also write the reflected equation for (33)

$$\partial_y S(Rz) = i\left\{\gamma,\sigma_3\right\}S(z) - i\sigma_3\partial_x S(Rz) - \xi S(Rz)\,. \tag{35}$$

Taking the derivative of (33) with respect to y and using (33), (35) and the Helmholtz equation we get

$$\left(m^2 + \left\{\gamma,\sigma_3\right\}^2 - \xi^2\right)S(z) = -i\left\{\xi,\sigma_3\right\}\partial_x S(z) - i\left\{\left\{\gamma,\sigma_3\right\},\xi\right\}S(Rz)\,. \tag{36}$$

The structure of singularities in this equation is different for the three terms, which have to cancel independently. This leads to

$$\left\{\xi,\sigma_3\right\} = 0\,, \tag{37}$$

$$\left\{\left\{\gamma,\sigma_3\right\},\xi\right\} = 0\,, \tag{38}$$

$$\left(m^2 + \left\{\gamma,\sigma_3\right\}^2 - \xi^2\right) = 0\,. \tag{39}$$

The same line of reasoning followed above to treat the translation symmetry in the y direction can be used to exploit the rotational symmetry of the Helmholtz equation. Considering rotations around L_2 with operator $(\partial_\theta - L_2\partial_y)$ where $\partial_\theta = x\partial_y - y\partial_x$, we have that $\left((\partial_\theta - L_2\partial_y) + (\partial_{\theta'} - L_2\partial_{y'})\right)G(z,z')$ is not singular for $z = z'$. Analyzing the singular terms we get the equation

$$\left((\partial_\theta - L_2\partial_y) + (\partial_{\theta'} - L_2\partial_{y'})\right)G(z,z') = -iL\,F^T(z)\sigma_3 S(z')\,. \tag{40}$$

Taking the derivative with respect to L_2, and following the same steps as before we arrive at

$$(\partial_\theta - L_2\partial_y)\,S(Rz) = -iqS(Rz) + iL\gamma\sigma_3 S(z)\,, \tag{41}$$

where

$$q = \begin{pmatrix} -a & \\ & 1-a \end{pmatrix}\,. \tag{42}$$

The reflected equation follows from replacing $z \to Rz$ and $(\partial_\theta - L_2\partial_y) \to -(\partial_\theta - L_1\partial_y)$,

$$(\partial_\theta - L_1\partial_y)\,S(z) = iqS(z) - iL\gamma\sigma_3 S(Rz)\,. \tag{43}$$

We derive (41) with respect to L_1, and use (33) to bring it to the form of the equation (43). From the comparison we obtain the following matrix differential equation

$$\xi = \frac{i}{L}\left(L\gamma^{-1}\frac{d\gamma}{dL}\sigma_3 + \gamma^{-1}q\gamma + q + \sigma_3\right)\,. \tag{44}$$

2.2 Parametrization

The algebraic equations (24), (34), (37 - 39) for the matrices are solved using the parametrization

$$\gamma = \frac{m}{2}\begin{pmatrix} u & b \\ c & u \end{pmatrix} \qquad ; \qquad \xi = m\begin{pmatrix} 0 & i\beta_1 \\ -i\beta_2 & 0 \end{pmatrix}, \tag{45}$$

where u, b, c, β_1, and β_2 are real functions of $t = mL$ and

$$u^2 + 1 = \beta_1\beta_2\,. \tag{46}$$

The differential equation (44) writes in components

$$\begin{aligned} \beta_1 &= \frac{tbu' - tb'u - ub}{t(u^2 - bc)}\,, & (47)\\ \beta_2 &= \frac{tcu' - tc'u - uc}{t(u^2 - bc)}\,, & (48)\\ 0 &= (1-2a)u^2 + tuu' - tc'b + 2(a-1)bc\,, & (49)\\ 0 &= (1-2a)u^2 - tuu' + tcb' + 2a\,bc\,. & (50) \end{aligned}$$

Defining $h = bc$ the last two equations give

$$h + \frac{t}{2}h' = tuu'\,. \tag{51}$$

Using (49) and (50) we eliminate $c'b$ and cb' in terms of u and h. Then (46) gives h in terms of u and using (51) it follows that

$$u'' + \frac{1}{t}u' - \frac{u}{1+u^2}u'^2 - u(1+u^2) - \frac{4u\left(a-\frac{1}{2}\right)^2}{t^2(1+u^2)} = 0\,. \tag{52}$$

This nonlinear ordinary differential equation can be transformed to take the form of a Painlevé V equation [12]. Taking into account that u must decay exponentially fast at infinity we also have

$$\int_t^\infty ds\; s\; u(s)^2 = \frac{t^2u'^2 - u^2\left((1-2a)^2 + t^2(1+u^2)\right)}{2(1+u^2)}\,. \tag{53}$$

2.3 Integrated quantities

The equations (7) and (20) give for the partition function

$$\partial_L\partial_{m^2}\log Z_a = -\int S^\dagger A\; S = 8\pi a\,(1-a)\,H\,, \tag{54}$$

where

$$H(L) = \int dxdy\; S_1^*(z)S_2(z)\,. \tag{55}$$

To find this quantity, we use the information obtained in the preceding Section. We first define the following auxiliary integrals

$$\begin{aligned} B_1(L) &= \int dxdy\; S_1^*(z)S_1(Rz)\,, & (56)\\ B_2(L) &= \int dxdy\; S_2^*(z)S_2(Rz)\,, & (57)\\ B_{12}(L) &= \int dxdy\; S_2^*(z)S_1(Rz)\,, & (58)\\ X_1(L) &= \int dxdy\; S_1^*(z)S_1(z)\,, & (59)\\ X_2(L) &= \int dxdy\; S_2^*(z)S_2(z)\,. & (60) \end{aligned}$$

These are convergent. They are also real since the relation (12) implies that $S_i^*(z) = S_i(\bar{z})$. Combining (33) and (35) we obtain

$$-i\left(\partial_y - i\partial_x\right)\left(S_1 S_2^*\right) = mu(S_1^R S_2^* + S_1 S_2^{*R}) - m\left(\beta_1 S_2 S_2^* + \beta_2 S_1 S_1^*\right) . \tag{61}$$

Integrating this equation in the plane it is

$$-i\int dxdy\left(\partial_y - i\partial_x\right)\left(S_1 S_2^*\right) = \frac{1}{8\pi a\left(1-a\right)} = 2muB_{12} - m\left(\beta_1 X_2 + \beta_2 X_1\right) , \tag{62}$$

where we have taken into account the pole of $S_1 S_2^*$ at L_1

$$S_1 S_2^* \sim \left(\left(4\pi\right)^2 a\left(1-a\right)\right)^{-1}\left(z - L_1\right)^{-1} , \tag{63}$$

and the fact that $S_1 S_2^*$ and its derivatives are continuous along the cut. Similarly, several other equations for the integrated variables can be constructed. From the angular equations (41) and (43) we get

$$0 = i\int dxdy\left(\partial_\theta - L_1\partial_y\right)\left(S_1 S_2^*\right) = tuB_{12} - \frac{t}{2}(bB_2 + cB_1) + H . \tag{64}$$

Using the eqs. (33) and (35) we obtain an expression for $\int dxdy\, S_2^{R*}\partial_y S_1$ in terms of B_1 and B_2, and a different expression for the same quantity arise using the angular equations (41) and (43). Combining them we get

$$0 = \frac{2}{t}(1-2a)B_{12} - cX_1 + bX_2 - \beta_1 B_2 + \beta_2 B_1 . \tag{65}$$

We still need three more equations involving the integral variables to completely determine them. A new one results as follows. It is easy to show that

$$\int S_1^* x\partial_x S_1 = -\frac{X_1}{2} = \int S_1^* y\partial_y S_1 , \tag{66}$$

$$\int S_2^* x\partial_x S_2 = -\frac{X_2}{2} = \int S_2^* y\partial_y S_2 . \tag{67}$$

Using the angular equations we have

$$\int S_1 y S_2^* = -i\int S_2^* x S_1 , \tag{68}$$

and from the eqs. (33) and (35), using (66) and (67),

$$\int S_1^* y\partial_x S_1 = -\int S_1^* x\partial_y S_1 = i\frac{X_1}{2} + \beta_1 m\int S_1^* y S_2 , \tag{69}$$

$$\int S_2^* y\partial_x S_2 = -\int S_2^* x\partial_y S_2 = -i\frac{X_2}{2} + \beta_2 m\int S_2^* x S_1 . \tag{70}$$

Now, inserting back these relations into the angular equations and taking into account (68) we finally have

$$-(1-\alpha)X_1\beta_2 + \alpha X_2\beta_1 + \frac{t}{2}(c\beta_1 - b\beta_2)B_{12} = 0 . \tag{71}$$

Additional linear equations for the integral variables can be obtained from the previous ones and the following ones for the derivatives,

$$H' = -(\frac{c}{2}B_1 + uB_{12} + \frac{b}{2}B_2) , \tag{72}$$

$$X_1' = -(uB_1 + bB_{12}) , \tag{73}$$

$$X_2' = -(cB_{12} + uB_2) . \tag{74}$$

These follow taking the derivative of (55), (59) and (60) with respect to L_2. Therefore all the integral variables can be written in terms of the functions u, b and c by solving a linear system. In particular we get

$$H = \left(16\pi a\left(1-a\right)m\right)^{-1} tu^2 . \tag{75}$$

2.4 Equations for the partition function and identification with Sine-Gordon model correlators

Combining (54) and (75) we have

$$L\partial_L \log Z_a = -\int_t^\infty dy\ y\ u^2(y)\,. \tag{76}$$

Comparing (52) and (76) with the results of [13], we see that the partition function satisfies the same equations than the inverse of a correlator of exponential operators in the SG model at the free fermion point. In fact it is

$$Z_a \simeq \frac{1}{\langle V_a V_{1-a}\rangle}\,, \tag{77}$$

where the correlator in the right hand side is evaluated in the Sine-Gordon model with action

$$\mathcal{A} = \int d^2x \left[\frac{1}{2}\partial_\nu\partial^\nu\phi - \mu\cos(\sqrt{4\pi}\phi)\right]\,, \tag{78}$$

and V_a is the operator $:\exp[i\sqrt{4\pi}a\phi]:$. From this identification we learn that the long distance behaviour of u is

$$u(t) \to \frac{2}{\pi}\sin(a\pi)K_{1-2a}(t)\,. \tag{79}$$

A correspondence between correlators of disorder operators for a scalar field with the inverse of correlators of exponential operators in the SG model was found in [11], by exploiting the equations for the form factors. Our results, obtained by a different method, agree with that calculation, when identifying our partition function Z_a with the correlator of boson disorder operators with charges a and $1-a$.

3 The entanglement and alpha entropies for free fields

Summarizing the results, the final expression for the α-entropies for integer $\alpha = n$ is

$$c_n = \frac{1}{2(1-n)}\sum_{k=1}^{n-1} w_{k/n}(t)\,, \tag{80}$$

with

$$w_a = = -\int_t^\infty dy\ y\ u_a^2(y)\,, \tag{81}$$

$$u_a^{''} + \frac{1}{t}u_a' = \frac{u_a}{1+u_a^2}\left(u_a'\right)^2 + u_a\left(1+u_a^2\right) + \frac{4(a-\frac{1}{2})^2}{t^2}\frac{u_a}{1+u_a^2}\,, \tag{82}$$

$$u_a(t) \to \frac{2}{\pi}\sin(a\pi)K_{1-2a}(t) \quad \text{as } t\to\infty\,. \tag{83}$$

We have introduced the $1/2$ factor in (80) because we want to present the results for a real scalar instead of a complex one. The function c_n is plotted in Figure 1 for some values of n.

In the conformal limit $t \to 0$ the function u diverges and has a series solution of the form

$$u_a(t) \sim \frac{1}{t\log(tb)} + a(a-1)t\log(tb) + \mathcal{O}(t)\,. \tag{84}$$

The full series is determined by the value of the integration constant b (which is a function of a). However, this is not known at present (except for the case $a = 1/2$ [10]). From eq. (53) we get for the leading terms

$$w_a(t) \sim -2a\,(1-a) - \frac{1}{\log(t)} + \mathcal{O}\left(\log^{-2}(t)\right)\,, \tag{85}$$



Figure 1: The c_n function for a real scalar field with, from top to bottom, $n = 2$, 3 and 50. At the origin they take the value $\frac{1+n}{6n}$ and they decay exponentially fast for large t.

and by analytic continuation of the sum in (80)

$$c_\alpha(t) = \frac{1+\alpha}{6\alpha} + \frac{1}{2\log(t)} + \mathcal{O}\left(\log^{-2}(t)\right) , \tag{86}$$

$$c(t) = \frac{1}{3} + \frac{1}{2\log(t)} + \mathcal{O}\left(\log^{-2}(t)\right) . \tag{87}$$

The first term corresponds to the conformal case for which there is a general result identifying $c = C/3$, where C is the Virasoro central charge [5]. Here it is $C = 1$ since we are considering a real scalar field.

The leading term for large distances of the entropic c-function can be obtained from (83) using the trick shown in [9]. We get

$$c(t) \sim \frac{1}{4}\, t\, K_1(2t) . \tag{88}$$

The analogous results for a Dirac field have been presented in a previous paper [9]. We copy here the corresponding formulae to facilitate the comparison with the bosonic case

$$c_n = \frac{1}{(1-n)} \sum_{k=-(n-1)/2}^{(n-1)/2} w_{k/n}(t) , \tag{89}$$

$$w_a = L\partial_L \log Z_a = -\int_t^\infty dy\; y\; u_a^2 , \tag{90}$$

$$u_a'' + \frac{1}{t}u_a' = -\frac{u_a}{1-u_a^2}\left(u_a'\right)^2 + u_a\left(1-u_a^2\right) + \frac{4a^2}{t^2}\frac{u_a}{1-u_a^2} , \tag{91}$$

$$u_a \to \frac{2}{\pi}\sin(\pi a) K_{2a}(t) \quad \text{as} \quad t \to \infty . \tag{92}$$

The similarities are striking. The reason is that here there is also an identification with SG correlators given by

$$Z_a \simeq \langle V_a V_{-a} \rangle . \tag{93}$$

The short and long distance behaviour of the Dirac c-function are

$$c(t) \sim \frac{1}{3} - \frac{1}{3}t^2\log^2(t) + \mathcal{O}\left(t^2\log(t)\right) \qquad \text{for} \quad t \to 0 , \tag{94}$$

$$c(t) \sim \frac{1}{2} t\, K_1(2t) \qquad \text{for} \quad t \to \infty . \tag{95}$$

We see that the bosonic c-function interpolates between the ones corresponding to a Dirac field (which has the same central charge at the conformal point) and a Majorana one (which has half the Dirac field



Figure 2: The entropic c-function $c(t) = LdS(L)/dL$ for a real scalar field (dotted line) and the Dirac and Majorana fermion ones (top and bottom dashed lines respectively). The bosonic c-function interpolates between the Dirac one at the origin, where both tend to the conformal value $c = 1/3$, and the Majorana c-function for large t, where they decay exponentially fast. The cusp at the origin of the bosonic function is due to a $1/\log(t)$ term. These curves were obtained by numerical computation on a lattice of up to 600 points.

entropy). However, it is evident that a deeper understanding of this phenomenon requires more than the explicit calculation presented here. We have plotted the entropic c-functions for bosons and fermions in Figure 2. This curves are calculated numerically putting the model in a lattice and using the method given in Appendix A. Note also the very different behaviour at the origin. The bosonic c-function has a $1/\log(t)$ term that can be ascribed to the zero mode which is present at the conformal point. In fact, for $t \to 0$ this behaviour gives for the entropy

$$S \sim \frac{1}{3}\log(L/\epsilon) + \frac{1}{2}\log\left(-\log(m\,\epsilon)\right) - \frac{1}{2}\log\left(-\log(m\,L)\right)\,, \tag{96}$$

where ϵ is an ultraviolet cutoff. In addition to the well-known logarithmic ultraviolet divergence, this last expression shows an explicit infrared divergence.

4 Universal terms for the entanglement entropy in arbitrary dimensions

Some universal terms for the entanglement entropy in more than two dimensions can be obtained from the two dimensional results. Specifically, let us consider a d-dimensional box, where d is the space dimension. Take one side of length L lying along the x_d axes direction to be much shorter than all the other sides having lengths R_i, $i = 1, ..., d-1$, and lying parallel to the x_i axes directions. The entanglement entropy of the box is divergent with a term proportional to the $(d-1)^{\text{th}}$-power of the cutoff and the total surface area [1, 2, 4]. Several other divergent terms proportional to lower dimensional geometric quantities also appear in the entropy, including a term with universal coefficient, proportional to the logarithm of the cutoff, which is due to the sharp angles at the box vertices [14]. However, one expects to find also subleading contributions which are finite and universal. In particular in the limit $R_i \to \infty$ one expects a leading universal term having the form

$$S \sim \text{non-univ} - k\frac{A}{L^{d-1}}\,, \tag{97}$$

where the area $A = \Pi_i R_i$, and k is a universal and dimensionless function of L and the renormalized parameters of the theory alone. For free bosons and fermions we can calculate exactly the function k. In the limit of large R_i we can compactify the x_i directions, with $i = 1, ..., d-1$, by imposing periodic boundary conditions $x_i = x_i + R_i$. As the term we are looking for is an extensive quantity in these coordinates, the compactification cannot change the result. For a real scalar field of mass m we Fourier

decompose it into the corresponding field modes in the compact directions. The problem then reduces to a one dimensional one with an infinity tower of massive scalar fields $\phi_{q_1,\ldots,q_{d-1}}$, where q_1,...,q_{d-1} are integers. These have masses given by

$$M^2_{q_1,\ldots,q_{d-1}} = m^2 + \sum_{i=1}^{d-1} \left(\frac{2\pi}{R_i} q_i\right)^2 . \tag{98}$$

Thus, summing over all the decoupled one dimensional fields, one has

$$L\frac{dS(L)}{dL} = \sum c\left(LM_{q_1,\ldots,q_{d-1}}\right) , \tag{99}$$

with $c(t)$ the one dimensional bosonic c-function. As we consider the large R_i limit we convert the sums into integrals and obtain

$$k(t) = \frac{\mathrm{vol}(\mathcal{S}^{d-2})}{(2\pi)^{d-1}}\, t^{d-1} \int_t^\infty dy_1\; y_1^{-d} \int_0^\infty dy_2\; y_2^{d-2}\; c\left(\sqrt{y_1^2+y_2^2}\right) , \tag{100}$$

where $\mathrm{vol}(\mathcal{S}^{d-2})$ is the volume of the $(d-2)$ dimensional unit sphere ($\mathrm{vol}(\mathcal{S}^0) = 2$). The same expression holds for free fermions. In this case the one dimensional c-function corresponds to the one of a Majorana field, and there is an additional factor of the dimension of the spinor space.

In particular, in the massless limit and in $2+1$ dimensions we have

$$k_S = \frac{1}{\pi}\int_0^\infty dt\; c_S(t) \simeq 0.039\,, \tag{101}$$

$$k_D = \frac{1}{\pi}\int_0^\infty dt\; c_D(t) \simeq 0.072\,, \tag{102}$$

for a real scalar (S) and a Dirac fermion (D) respectively. The values on the right hand side were found by numerical integration of the curves of Figure 2. We have checked these last results numerically on a two dimensional lattice for rectangular sets of size up to 10×100 points, finding a perfect accord within a 3% error. Further details of this and other results on universal terms for the entropy in $2+1$ dimensions will be published elsewhere [14].

5 Conclusions

In this work we have completed the discussion on the alpha and entanglement entropies for free massive fields in two dimensions, presenting the results corresponding to a scalar field. The Dirac fermion case was analyzed in a previous paper by a different approach. Here we arrive to the exact expression for the α-entropies for integer α and expansions for short and long distances of the geometric entropy for a massive scalar field. We do it by direct calculation of the relevant information on the Green function on a plane, where boundary conditions are imposed on a finite interval. In both cases, bosonic and fermionic, we find a relation to correlators of exponential operators in the Sine-Gordon model, associated to the same type of non linear differential equation of the Painlevé V type.

As we previously noted for the fermionic case [9], also for the scalar field the expression for the c_α functions show that they are decreasing in an explicit manner (see (81) and (90)). This could be a hint to a possible extension of the entanglement entropy c-theorem.

The method used in this work seems to be very well suited to generalizations, and thus it might be useful to calculate different contributions to the geometric entropy in more dimensions. It essentially gives information from the translation and rotation symmetries of the wave equation, even in the presence of symmetry breaking boundary conditions, once one has understood the structure of the singularities.

Concerning the geometric entropy in arbitrary dimensions, we have also discussed universal terms which are related to the two dimensional results. We have found a universal term of the form

$$S_U \sim -k\frac{A}{L^{d-1}} \tag{103}$$

for the entanglement entropy of a thin box in d spatial dimensions, with a small side of length L and big phase area A. In some sense, this term could be interpreted as a universal (cutoff independent) form of the area law for the entropy, with L playing the role of the cutoff. It is known [7] that in quantum field theory a universal measure of information must be expressed in terms of the mutual information $I(X,Y) = S(X) + S(Y) - S(X \cup Y)$ between two different non intersecting sets X and Y, rather than the entropy for a single set. In this sense it is remarkable that the term (103) determines the mutual information between to big boxes as their parallel phases come into contact.

6 Acknowledgments

We thank Cesar Fosco for very useful discussions all along the realization of this project.

7 Appendix A: numerical entropy in a lattice

We use the method presented in [15] to give an expression for ρ_A in terms of correlators for free boson and fermion discrete systems. The case of a fermion field is summarized in [9]. For the scalar field it is a generalization of the method given by Sredniki [2]. We start with a free Hamiltonian for bosonic degrees of freedom with the form

$$H = \frac{1}{2}\sum \pi_i^2 + \frac{1}{2}\sum_{ij} \phi_i M_{ij} \phi_j \ , \tag{104}$$

where $\phi_i$and π_i obey the canonical commutation relations $[\phi_i, \pi_j] = i\delta_{ij}$, and M is a Hermitian positive definite matrix. The vacuum (ground state) correlators are given by

$$X_{ij} = \langle \phi_i \phi_j \rangle = \frac{1}{2}(M^{-\frac{1}{2}})_{ij} \, , \tag{105}$$

$$P_{ij} = \langle \pi_i \pi_j \rangle = \frac{1}{2}(M^{\frac{1}{2}})_{ij} \, . \tag{106}$$

Let X_{ij}^A and P_{ij}^A be the correlator matrices restricted to the region A, that is $i, j \in A$. Due to the Wick theorem any other correlator inside A can be written in terms of these two. The expectation values calculated with ρ_A must also satisfy Wick's theorem and give place to the two point correlators X_{ij}^A and P_{ij}^A. This fixes ρ_A to be of the form

$$\rho_A = K \, e^{-\Sigma \epsilon_l a_l^\dagger a_l} \, , \tag{107}$$

where K is a normalization constant and the a_l are independent boson annihilation operators, $[a_i, a_j^\dagger] = \delta_{ij}$, which can be expressed as linear combination of the ϕ_i and π_j, i, $j \in A$. The ϵ_l are related to the eigenvalues ν_l of $\sqrt{X^A.P^A}$ by

$$\tanh(\frac{\epsilon_l}{2}) = \frac{1}{2\nu_l} \, . \tag{108}$$

Hence, the entropy can be evaluated as $S_A = \Sigma \, S_l$, where

$$S_l = -\log(1 - e^{-\epsilon_l}) + \epsilon_l \frac{e^{-\epsilon_l}}{1 - e^{-\epsilon_l}} = (\nu_l + \frac{1}{2})\log(\nu_l + \frac{1}{2}) - (\nu_l - \frac{1}{2})\log(\nu_l - \frac{1}{2}) \, . \tag{109}$$

Thus, to compute the entropy numerically we need to diagonalize the matrix $X^A.P^A$. Its eigenvalues ν_l^2, due to the uncertainty relations, are always greater than $1/4$. Note that for the total system $XP = \frac{1}{4}$ and $\nu_l \equiv \frac{1}{2}$, with zero entropy.

The lattice Hamiltonian for a real massive scalar is given by

$$\mathcal{H} = \frac{1}{2}\sum_{n=0}^{N-1} \left(\pi_n^2 + (\phi_{n+1} - \phi_n)^2 + m^2\phi_n^2\right) \, . \tag{110}$$

We have taken N lattice sites and set the lattice spacing to one. The correlators (105) and (106) for the infinite lattice limit $N \to \infty$ are

$$\langle \phi_n \phi_m \rangle = \int_0^1 dx \frac{e^{2\pi i x(m-n)}}{2\sqrt{m^2 + 2(1-\cos(2\pi x))}}, \tag{111}$$

$$\langle \pi_n \pi_m \rangle = \int_0^1 dx \frac{1}{2} e^{2\pi i x(m-n)} \sqrt{m^2 + 2(1-\cos(2\pi x))}. \tag{112}$$

References

[1] L. Bombelli, R. K. Koul, J. H. Lee and R. D. Sorkin, Phys. Rev. D **34**, 373 (1986).

[2] M. Srednicki, Phys. Rev. Lett. **71**, 666 (1993) [arXiv:hep-th/9303048].

[3] C. G. Callan and F. Wilczek, Phys. Lett. B **333**, 55 (1994) [arXiv:hep-th/9401072].

[4] H. Casini, Class. Quant. Grav. **21**, 2351 (2004) [arXiv:hep-th/0312238].

[5] C. Holzhey, F. Larsen and F. Wilczek, Nucl. Phys. B **424**, 443 (1994) [arXiv:hep-th/9403108].

[6] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, Phys. Rev. Lett. **90**, 227902 (2003) [arXiv:quant-ph/0211074]; J. I. Latorre, E. Rico and G. Vidal, Quant. Inf. and Comp. **4**, 048 (2004) [arXiv:quant-ph/0304098]; J. I. Latorre, C. A. Lutken, E. Rico and G. Vidal, Phys. Rev. A **71**, 034301 (2005) [arXiv:quant-ph/0404120]; R. Orus, Physical Review A **71**, 052327 (2005) [arXiv:quant-ph/0501110]; V.Vedral, [arXiv:quant-ph/0410021]; M. B. Plenio, J. Eisert, J. Dreissig and M. Cramer, Phys. Rev. Lett. **94**, 060503 (2005) [arXiv:quant-ph/0405142]; P. Calabrese and J. Cardy, JSTAT **0406**, P002 (2004) [arXiv:hep-th/0405152]; A. Osterloh, L. Amico, G. Falci, R. Fazio, Nature **416**, 608 (2002) [arXiv:quant-ph/0202029]; F. Verstraete, M. A. Martin-Delgado, J. I. Cirac, Phys. Rev. Lett. **92**, 087201 (2004) [arXiv: quant-ph/0311087]; A. R. Its, B.-Q. Jin, V. E. Korepin, Journal Phys. A: Math. Gen. **38**, 2975 (2005) [arXiv: quant-ph/0409027]; V. Korepin, Phys. Rev. Lett. **92**, 096402 (2004) [arXiv: cond-mat/0311056]; J. P. Keating, F. Mezzadri, Phys. Rev. Lett. **94**, 050501 (2005) [arXiv: quant-ph/0504179]; E. Rico Ortega, Ph.D. Thesis [arXiv:quant-ph/0509037].

[7] H. Casini and M. Huerta, Phys. Lett. B **600**, 142 (2004) [arXiv:hep-th/0405111].

[8] A. Kitaev and J. Preskill [arXiv:hep-th/0510092].

[9] H. Casini, C. D. Fosco and M. Huerta, JSTAT **P07007** (2005) [arXiv:cond-mat/0505563].

[10] J. M. Myers, J. Math. Phys., **6**, 1839 (1965).

[11] G. Delfino, P. Grinza and G. Mussardo, Phys. Lett. B **536**, 169 (2002) [arXiv:hep-th/0203063].

[12] E. L. Ince, *Ordinary differntial equations*, Dover Publications Inc., USA, 1926.

[13] D. Bernard and A. LeClair, Nucl. Phys. B **426**, 534 (1994) [Erratum-ibid. B **498**, 619 (1997)] [arXiv:hep-th/9402144]; H. Widom, J. Int. Eqs. Appl. **10**, 363 (1998) [arXiv: solv-int/9702007].

[14] H. Casini, C. D. Fosco and M. Huerta, in preparation.

[15] I. Peschel, J. Phys. A: Math. Gen. **36**, L205 (2003) [arXiv:cond-mat/0212631]; M. C. Chung, I. Peschel, Phys. Rev. B **62**, 4191 (2000) [arXiv:cond-mat/0004222].